

Patient Education - Oxymetazoline 1.2%, Hyaluronic Acid 0.5% Topical Gel

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes.

Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Oxymetazoline is a decongestant that is used to treat facial redness caused by rosacea, it works by shrinking dilated blood vessels. Oxymetazoline topical may also be used for purposes not listed in this medication guide.

To make sure oxymetazoline topical is safe for you, tell your doctor if you have: heart disease, high blood pressure blood circulation problems glaucoma or peripheral vascular disease such as Raynaud's syndrome, Buerger's disease, scleroderma, or Sjögren's syndrome.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether oxymetazoline topical passes into breast milk or if it could affect the nursing baby. Tell your doctor if you are breast-feeding.

Oxymetazoline is not approved for use by anyone younger than 18 years old.

Do not take by mouth. Topical medicine is for use only on the skin. Do not use this medicine on open wounds or irritated skin. If this medicine gets in your eyes, nose, mouth, or vagina, rinse with water.

Wash your hands after applying this medicine.

Call your doctor if your symptoms do not improve, or if they get worse.

Store at room temperature away from moisture and heat.

Oxymetazoline topical side effects:

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Stop using this medicine and call your doctor at once if you have: worsening of your rosacea symptoms numbness, tingling, cold feeling in your hands or feet pale or purple appearance in your fingers or toes or blurred vision, tunnel vision, eye pain, or seeing halos around lights.

Common side effects may include: skin redness or itching pain or other skin reactions.

This is not a complete list of side effects and others may occur.