

Patient Education - Desonide 0.04%, Idochlorhydroxyquin 3% Topical Gel

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

- Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website address: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Desonide - Topical Corticosteroid

Clioquinol - Antifungal

WHAT IS THIS DRUG USED FOR? It is used to treat infected/irritated skin conditions.

WHAT DO I NEED TO TELL MY DOCTOR BEFORE I TAKE THIS DRUG? If you have an allergy to desonide, clioquinol or any other part of this drug. If you are allergic to this drug any part of this drug or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had. If there is an infection where this drug will be used. If you have thinning of the skin where you are putting this drug. This is not a list of all drugs or health problems that interact with this drug. Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

WHAT ARE SOME THINGS I NEED TO KNOW OR DO WHILE I TAKE THIS DRUG? Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists. Talk with your doctor before you use other drugs or products on your skin. Do not use tight-fitting diapers or plastic pants if treated part is in the diaper area. This may cause more drug to get into the body. Do not put on cuts, scrapes, or damaged skin. Use care when putting on a large part of the skin or where there are open wounds. Talk with the doctor. Do not use this drug for longer than you were told by your doctor. This drug may raise the chance of cataracts or glaucoma. Talk with the doctor. Use with care in children. Talk with the doctor. This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor. Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby. If you are breast-feeding and use this drug on your breast or nipple, wash the area before breast-feeding your child.

WHAT ARE SOME SIDE EFFECTS THAT I NEED TO CALL MY DOCTOR ABOUT RIGHT AWAY? WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash hives itching red, swollen, blistered, or peeling skin with or without fever wheezing tightness in the chest or throat trouble breathing, swallowing, or talking unusual hoarseness or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss. Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache, or slow healing. Skin irritation. Skin changes (pimples, stretch marks, slow healing, hair growth). Change in color of skin. Change in eyesight, eye pain, or very bad eye irritation.

WHAT ARE SOME OTHER SIDE EFFECTS OF THIS DRUG? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Burning. Stinging. Dry skin. Signs of a common cold. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

HOW IS THIS DRUG BEST TAKEN? Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely. Use as you have been told, even if your signs get better. Do not take this drug by mouth. Use on your skin only. Keep out of your mouth, nose, and eyes (may burn). Wash your hands before and after use. Do not wash your hands after use if putting this on your hand. Clean affected part before use. Make sure to dry well. Put a thin layer on the affected skin and rub in gently. Do not use coverings (bandages, dressings, make-up) unless told to do so by the doctor. Do not put on the face, underarms, or the groin area unless told to do so by the doctor.

HOW DO I STORE AND/OR THROW OUT THIS DRUG? Store at room temperature. Do not freeze. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.

GENERAL DRUG FACTS If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider. If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.