

Patient Education - Miconazole 0.5%, Cholestyramine 10% in 25% Zinc Oxide Topical Paste

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program

Website address: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Miconazole topical is an antifungal medication that prevents fungus from growing on your skin. It is used to treat skin infections such as athlete's foot, jock itch, ringworm, tinea versicolor (a fungus that discolors the skin), and yeast infections of the skin & may also be used for purposes not listed in this medication guide.

You should not use miconazole topical if you are allergic to miconazole.

Ask a doctor or pharmacist if it is safe for you to use this medicine if you have other medical conditions, especially if you are using a blood thinner such as warfarin, Coumadin, Jantoven.

It is not known whether miconazole topical will harm an unborn baby. Do not use this medicine without a doctor's advice if you are pregnant. It is not known whether miconazole topical passes into breast milk or if it could harm a nursing baby. Do not use this medicine without a doctor's advice if you are breast-feeding a baby.

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended.

Do not take by mouth. Miconazole topical is for use only on the skin. Do not use this medicine on open wounds.

Miconazole topical is not for use in the vagina or rectum.

Wash your hands before and after using this medication, unless you are treating a skin condition on your hands.

Do not cover the treated skin area unless your doctor tells you to. A light cotton-gauze dressing may be used to protect clothing.

Use this medication for the full prescribed length of time. Your symptoms may improve before the infection is completely cleared. Skipping doses may also increase your risk of further infection that is resistant to antifungal medicine.

Store at room temperature away from moisture and heat. Keep the tube tightly closed when not in use.

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat.

Stop using miconazole topical and call your doctor at once if you have: severe blistering, redness, or irritation of treated skin.

Common side effects may include: itching, peeling, or dry skin.

Although the risk of serious side effects is low when miconazole topical is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream, including:

dry mouth, sore tongue, tooth pain, red or swollen gums altered sense of taste nausea, diarrhea or headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

Cholestyramine is typically used to reduce cholesterol (fatty acids) in the blood, but in topical cream preparations it is used to decrease skin rash and irritation associated

with a dermatologic or other underlying condition.

Cholestyramine may also be used for purposes not listed in this medication guide. Generally, topical applications are not absorbed in significant amounts into the blood stream and systemic adverse effects are uncommon. Call your medical provider if you experience any unusual/severe gastrointestinal pain, constipation, blood in urine or stools (including black or tarry),

Zinc Oxide is used to treat skin irritation by creating a protective barrier. Side effects are uncommon. If you have any specific questions about Zinc Oxide, please contact the pharmacist or your physician.