

Patient Education - Ivermectin 1.2%, Metronidazole 1.2%, Niacinamide 2% Topical Gel

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes. Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Ivermectin - used to treat rosacea.

Wash your hands before and after use.

Do not take ivermectin cream by mouth. Use on your skin only. Keep out of your mouth, nose, and eyes (may burn). Avoid getting ivermectin cream in the eyes or on the lips.

Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

Signs of an allergic reaction, like rash hives itching red, swollen, blistered, or peeling skin with or without fever wheezing tightness in the chest or throat trouble breathing, swallowing, or talking unusual hoarseness or swelling of the mouth, face, lips, tongue, or throat.
Very bad irritation where ivermectin cream is used. Very bad burning or burning that does not go away.

More common side effects include: Irritation where ivermectin cream is used. Burning.

Metronidazole is an antibiotic that fights bacteria in the body used to treat skin lesions caused by rosacea & may also be used for purposes not listed in this medication guide. You should not use metronidazole topical if you are allergic to it.

Metronidazole topical is not expected to harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant. It is not known whether metronidazole topical passes into breast milk or if it could harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Wash and gently dry your skin before applying metronidazole. Apply the medication in a thin layer and rub in completely. You may apply cosmetics after using metronidazole topical.

Store metronidazole topical at room temperature away from moisture or heat.

Do not take by mouth. Metronidazole topical is for use only on the skin. If this medication gets in your eyes, rinse with water. Do not use topical metronidazole in the vagina.

Get emergency medical help if you have signs of an allergic reaction: hives difficulty breathing swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have severe stinging or burning when you apply metronidazole topical.

Common side effects may include: mild burning or stinging numbness or tingly feeling in your hands or feet mild skin redness or irritation dry, scaly, or itchy skin nausea or metallic taste in your mouth. This is not a complete list of side effects and others may occur.

Niacinamide – Topical anti-inflammatory

When applied topically, niacinamide has been shown to have anti-inflammatory effects on the skin. It has been used to treat skin conditions like acne and rosacea, a facial skin disorder characterized by redness.

Possible side effects include: mild burning, itching, or redness.