

Patient Education - Latanoprost 0.01%, Minoxidil 7.5%, Finasteride 0.25% Topical Solution

Hillcrest Pharmacy 781 E FT Union Blvd Midvale, UT 84047 – Direct 385-900-1400, Toll Free 1-855-499-2556

Patient Education Insert

-Apply medication to affected area as prescribed

This medication was compounded by a pharmacy per prescriber request. For external use only. Wash hands before and after use unless the medication is for your hands. Store at room temperature (68 to 77 degrees Fahrenheit). Keep away from children and pets. Do not flush unused medications or pour down a sink or drain. Do not swallow and avoid contact with eyes.

Do not use more than your medical provider has instructed you to use. Do not use more often or for longer than you were instructed. Doing any of these may increase the chance of undesirable side effects.

The “Do not use after” date printed on the prescription label is generated using State & Federal compounding laws. Some preparations are made 1-3 days before the prescription is issued, in anticipation of expected demand. If you have any questions or concerns regarding this, please do not hesitate to call and speak to one of our pharmacists.

Do not use if an allergic reaction occurs due to any of the listed active ingredients on the prescription label. Signs of an allergic reaction may include rash hives itching shortness of breath wheezing cough swelling of face, lips, tongue or throat or any other signs.

Contact your medical provider or pharmacy if you have concerns about listed side effects or if you experience unexpected irritation. Notify your medical provider of any medical updates and of all drug changes including over-the-counter and herbal medications.

If you miss an application, apply as soon as you remember. If it is close to the next application, skip the missed application. Do not double applications.

We want to offer you the best service possible. In the event you have a concern or complaint please call the pharmacy at 385-900-1400.

MedWatch: The FDA Safety Information and Adverse Event Reporting Program - Website: <https://www.fda.gov/Safety/MedWatch/default.htm> - Toll free number: 1-888-463-6332

Your medication contains the following active ingredients. Please review the following:

Latanoprost - is traditionally used to treat certain types of glaucoma and other causes of high pressure inside the eye, however, is being used off-label to treat hair-loss (alopecia) in this application and may also be used for purposes not listed in this medication guide.

You should not use latanoprost ophthalmic if you are allergic to latanoprost or benzalkonium chloride.

Latanoprost ophthalmic is not approved for use by anyone younger than 18 years old.

Avoid eye exposure for this type of latanoprost preparation. Ophthalmic exposure may cause a gradual change in the color of your eyes or eyelids and lashes, usually an increase in brown pigment. You may also notice increased growth or thickness of your eyelashes. These changes occur slowly and you may not notice them for months or years. Color changes may be permanent even after your treatment ends, and may occur only in the eye being treated. This could result in a cosmetic difference in eye or eyelash color from one eye to the other.

Tell your doctor if you are pregnant or breastfeeding as exposure may result in serious side effects to a fetus or nursing infant.

Get emergency medical help if you have signs of an allergic reaction: hives difficult breathing swelling of your face, lips, tongue, or throat. This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

Minoxidil - used to help you regrow hair on your scalp.

Minoxidil topical will not cause permanent regrowth of scalp hair. You must continue using the product to keep the regrowth of your hair and may also be used for purposes not listed in this medication guide.

Do not use minoxidil topical if the skin on your scalp is red, swollen, irritated, or infected. This may allow more of the medication to be absorbed by your body, which could be dangerous.

Check the medicine label carefully to make sure minoxidil topical will treat your specific type of hair loss.

Using more of this medicine than recommended will not speed up hair growth, and may cause dangerous side effects.

Ask a doctor or pharmacist if it is safe for you to use minoxidil topical if you have other medical conditions, especially heart disease.

FDA pregnancy category C. It is not known whether minoxidil topical will harm an unborn baby. Do not use this medicine without a doctor's advice if you are pregnant. It is not known whether minoxidil topical passes into breast milk or if it could harm a nursing baby. Do not use this medicine without a doctor's advice if you are breast-feeding a baby.

Do not use this medicine on anyone under 18 years old without medical advice.

Although the risk of serious side effects is low when minoxidil topical is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Call your doctor at once if you have: severe scalp irritation unwanted growth of facial hair chest pain, fast heartbeats swelling in your hands or feet, rapid weight gain a light-headed feeling, like you might pass out headache, dizziness, confusion or flushing (warmth, redness, or tingly feeling).

Common side effects may include: changes in the color or texture of your hair.

Finasteride - A 5 Alpha-Reductase Inhibitor, used to help you regrow hair on your scalp by preventing the conversion of testosterone to dihydrotestosterone (DHT) in the body. Finasteride may also be used for purposes not listed in this medication guide

Finasteride can cause birth defects if a woman is exposed to it during pregnancy & should not be handled by a woman who is pregnant or may become pregnant. If a woman accidentally comes into contact with finasteride, wash the area with soap and water right away.

You should not use finasteride if you are allergic to it. Finasteride should never be taken by a woman or a child. Finasteride can be absorbed through the skin, and women or children should not be permitted to handle finasteride tablets.

Using finasteride may increase your risk of developing prostate cancer. Talk with your doctor if you have concerns about this risk. Tell your doctor if you have ever had: liver disease, or abnormal liver function tests, prostate cancer, a bladder muscle disorder, stricture of your urethra urination problems or an allergic reaction to a similar medicine called dutasteride (Avodart).

Get emergency medical help if you have signs of an allergic reaction: hives difficulty breathing swelling of your face, lips, tongue, or throat.

Call your doctor at once if you notice any signs of male breast cancer, such as: breast lumps, breast pain or tenderness, nipple discharge or any other breast changes.

Common side effects may include: loss of interest in sex impotence trouble having an orgasm or abnormal ejaculation.

The sexual side effects of finasteride may continue after you stop taking finasteride. Talk to your doctor if you have concerns about these side effects.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.